

PALYNOSTRATIGRAPHY AND PALAEOGEOGRAPHY OF EARLY PERMIAN STRATA IN THE ZAGROS BASIN, SOUTHEAST- SOUTHWEST IRAN

M. Ghavidel-syooki

*Exploration and Production Division, National Iranian Oil Company, P.O. Box: 1065,
Tehran, Islamic Republic of Iran*

Abstract

A total of 500 Early Permian samples from the Chal-i-Sheh Formation in the Zagros Basin were examined palynologically in order to determine the stratigraphical age of the formation and to assess the palaeogeographical relationships of this basin to the Southern and Northern Hemispheres during the Early Permian. Fifty-eight taxa were recorded including 46 pollen, 10 spore and 2 acritarch species, which permitted the establishment of the age of the formation. Quantitatively, pollen contributed 98 percent, spores 1.5 percent, and acritarchs 0.5 percent of the total assemblages. The high relative percentage of gymnospermous pollen taxa suggests a relatively cold condition which was in favor of gymnospermous plants at the beginning of the Early Permian. Comparison of the taxa recorded from the Early Permian strata of the Zagros Basin with those reported from other parts of the world indicates that the Zagros Basin was part of the Gondwanan Supercontinent located on the southern shore of the Tethys sea during the Permian period.

Introduction

During the past decade, the writer has carried out a detailed palynological study on the Palaeozoic sequences (Barut, Zaigun, Lalun, Mila, Ilebeck, Zard-Kuh, Seyahou, Sarchahan and "Faraghan" Formations) in the Zagros Basin (Fig. 2). The Zagros Basin includes the southeastern part of Iran and most of the adjacent Persian Gulf, forming a SE-NW trending linear belt, approximately 1400 km long and 250 km wide, which is separated from the Central Iranian Basin by the 5-10 km wide Main Zagros Thrust or Crush Zone (Fig. 1).

Among the Palaeozoic rock units of this basin, the

"Faraghan" Formation has received minimal biostratigraphic interest since it lacks marine fauna. However, based on its stratigraphic position and some plant remains, it has been assigned to the Early Permian [62] or the Permo-Carboniferous period [57]. The writer's palynological investigations have, however, revealed that part of the "Faraghan" Formation was deposited in the Devonian period and another part during the Early Permian [26]. Therefore, the carboniferous strata are apparently unrepresented in the Zagros Basin.

As a result of these studies, the use of the term "Faraghan" Formation was discontinued and, with the approval of the National Iranian Stratigraphic Committee, was replaced with the terms Zakeen Formation (of

Keywords: Chal-i-Sheh formation; Early Permian; Miospores; Palaeogeography; Zagros Basin

Devonian age) and Chal-i-Sheh Formation (of Permian age). This paper aims to demonstrate some diagnostic palynomorph taxa from the Chal-i-Sheh Formation in order to resolve aspects of palaeogeographic relationships of the Zagros Basin during the Early Permian. The palaeogeography of the Zagros Basin is important at this geological time since the palynological data are able to make clear the relationships of this basin to central and northern parts of Iran as well as to other parts of the world.

Previous Studies

The Zagros Basin encompasses the southeastern part of Iran and most of the Persian Gulf and has been a major center of oil exploration. Most of the formation nomenclature for the sedimentary sequence of the Zagros Basin was established by petroleum geologists (J.V. Harrison, 1930, unpublished company report). Setudehnia [57] described the Palaeozoic strata in parts of the Zagros Basin such as Zard-Kuh, Kuh-e-Gereh, Kuh-e-Dinar and Oshtoran-Kuh areas (Fig. 1). As a result of field studies, the Pre-Triassic sediments of the Zagros Basin were named by Szabo and Kheradpir [62] and their rock units approved by the National Iranian Stratigraphic Committee in that year. Although Szabo and Kheradpir's paper is the most complete documentation for the Palaeozoic rock units in the Zagros Basin, the age assignments of some of them are still incomplete or in premature stages due to a lack of diagnostic biostratigraphical evidence. With the exception of the Zakeen and Chal-i-Sheh Formations, which lack marine fauna, the rest of the Palaeozoic rock units in the basin contain well-preserved and abundant faunas consisting of trilobite, coral, brachiopod, graptolite and fusulinid, providing reliable age controls. The studied sequence was originally referred to as Carboniferous Sandstones or Permo-Carboniferous Sandstones by Setudehnia [57], but later was named the Faraghan Formation by Szabo and Kheradpir [62] and assigned to the Lower Permian. The age assignment of the sequence either by Setudehnia [57], or Szabo and Kheradpir [62], was based on Seward's work [58] which identified plant remains as *Sigillaria persica* and suggested an Upper Carboniferous (Stephanian), or Lower Permian age.

The writer's preliminary research focused on palynological characteristics of the sequence (Faraghan Formation) in the Gahkum-Faraghan areas (Fig. 1). The study revealed that 200 out of 245 m of the Faraghan Formation in both Kuh-e-Faraghan and Kuh-e-Gahkum have been deposited during the Devonian period and the remainder laid down during the Early Permian [26]. This resulted in the subdivision of the Faraghan Formation into the Zakeen Formation (Devonian) and the Chal-i-Sheh Formation (Early Permian) by the National Iranian Stratigraphic Committee in 1996. This paper is concerned

with the palynological characteristics of the Chal-i-Sheh Formation (Fig. 2) and those of the Zakeen formation will be published in another paper.

Stratigraphy

The name Chal-i-Sheh Formation is given to the Early Permian part of the succession formerly known as the Faraghan Formation. This rock unit is well-developed in all of the high mountain ranges in the Zagros Basin, from northwest to southeast parts of Iran (Fig. 1).

The Chal-i-Sheh Formation was named as one of the best developed outcrop sections in the Chal-e-Sheh area [30]. Its thickness varies from a minimum of a few meters at Kuh-e-Dinar to 500 meters in the type locality area (Fig. 2). The formation mainly consists of sandstone, siltstone, shale and a few limestone beds.

The Chal-i-Sheh Formation lacks marine fauna but contains abundant plant remains at one horizon in the type locality area (Fig. 2). The plant remains were collected by J.V. Harrison in 1930 and identified by Seward [58] as *Sigillaria persica* suggesting a Late Carboniferous (Stephanian), or Early Permian age. The upper contact of the Chal-i-Sheh Formation is gradational with the Dalan Formation (Upper Permian) at all surface and subsurface sections (Fig. 2) and is marked by a transition from interbedded sandstones to limestone beds. The Lower contact is characterized by a major hiatus, the magnitude of which increases in a SE-NW direction in the Zagros Basin with the lowest beds of the Chal-i-Sheh Formation resting on a variety of older strata (Fig. 2).

Laboratory Techniques

A total of 500 surface and subsurface samples from the Chal-i-Sheh Formation were selected for palynological study. Digestion of the samples was carried out using standardized techniques. All slides are housed in the Palaeontological Section of the Exploration Department of the National Iranian Oil Company.

List of Taxa Identified

In this study, no attempt was made to establish systematic description for the known morphotypes of the Chal-i-Sheh Formation based on conventional suprageneric classification, instead, they were arranged under their probable parent plant groups. This informal scheme has already been used for Australian morphotypes by Gilby and Foster [31] who attributed the Permian morphotypes to the categories of articulates, pterophytes, lycophytes and gymnosperms. In the list of taxa given below, plate and figure numbers given in brackets refer to the illustrations in this paper.




Figure 1. Distribution of the Chal-i-Sheh Formation in the Zagros Basin and the locations of the study area to the main Zagros Thrust, Southern Iran




Figure 2. Correlation chart of Chal-i-Sheh Formation and its relationship to overlying and underlying formation in the Zagros Basin of Iran

Articulates

Punctatisporites gretensis, Balme and Hennelly, 1965 (Pl. I, Fig. 5).

Pterophytes

Horriditriletes ramosus, (Balme and Hennelly) Bharadwaj and Salujha, 1964 (Pl. I, Figs. 2 & 6).

Laevigatosporites colliensis, (Bharadwaj and Hennelly) Venkatachala and Kar, 1968 (Pl. I, Fig. 10).

Leiotriletes virkkii, Tiwari, 1965 (Pl. I, Fig. 9).

Lophotriletes scotinus, Segroves, 1970 (Pl. I, Fig. 1).

Pseudoreticulatispora pseudoreticulata, (Balme and Hennelly) Bharadwaj and Srivastava 1969 (Pl. I, Fig. 4).

Verrucosiporites andersonii, (Anderson) Backhouse, 1988 (Pl. I, Fig. 8).

Lycopods

Densoisporites solidus, Segroves, 1970 (Pl. I, Fig. 7).

Jayantisporites variabilis, (Anderson) Backhouse, 1991 (Pl. I, Fig. 3).

Kraeuselisporites niger, Segroves, 1970 (Pl. I, Fig. 12).

Gymnospermous Pollen**Monosaccate Pollen**

Bascanisporites undosus, Balme and Hennelly, 1956 (Pl. III, Fig. 4).

Nuskoisporites triangularis, Potonié and Lele, 1961 (Pl. II, Fig. 7).

Plicatipollenites densus, Srivastava, 1970 (Pl. II, Fig. 9).

Plicatipollenites indicus, Lele, 1964 (Pl. V, Fig. 12).

Potoniéisporites granulatus, Bose and Kar, 1966 (Pl. II, Fig. 12).

Potoniéisporites neglectus, Potonié and Lele, 1961 (Pl. II, Fig. 6).

Rugasaccites orbiculatus, Lele and Maithy, 1969 (Pl. II, Fig. 5).

Disaccate Non-Taeniate

Alisporites nuthallensis, Clarke, 1965 (Pl. II, Fig. 2).

Caheniasaccites ellipticus, Bose and Maheshwari, 1968 (Pl. II, Fig. 1).

Caheniasaccites indicus, Srivastava, 1970 (Pl. II, Fig. 10).

Falcisporites, sp., (Pl. II, Fig. 8).

Klausipollenites schaubergeri (Potonié and Klaus) Jansonius, 1962 (Pl. II, Fig. 11).

Walikalesaccites ellipticus, Bose and Kar, 1966 (Pl. V, Fig. 2).

Colpate Pollen

Cycadopites cymbatus, (Balme and Hennelly) Segroves, 1969 (Pl. III, Fig. 6).

Praecolpatites sinuosus, (Balme and Hennelly) Bharadwaj and Srivastava, 1969 (Pl. II, Fig. 3).

Taeniate Pollen

Boutakoffites elongatus, Bose and Kar, 1966 (Pl. IV, Fig. 3).

Complexisporites polymorphus, Jizba, 1962 (Pl. VI, Figs. 1 & 12).

Corisaccites alutas, Venkatachala and Kar, 1966 (Pl. V, Fig. 10).

Costapollenites ellipticus, Tschudy and Kosanke, 1966 (Pl. IV, Fig. 6).

Crustaesporites globosus, Leschik, 1956 (Pl. III, Fig. 7).

Decussatisporites circularis, Sinha, 1970 (Pl. III, Fig. 10).

Distriamonocolpites ovalis, Bharadwaj and Sinha, 1969 (Pl. III, Fig. 11).

Fusacolpites fusus, Bose and Kar, 1966 (Pl. IV, Fig. 12).

Fusacolpites ovatus, Bose and Kar, 1966 (Pl. IV, Fig. 8).

Hamiapollenites karrooensis, (Hart) Hart, 1964 (Pl. VI, Fig. 7).

Hamiapollenites saccatus, Wilson, 1962 (Pl. V, Fig. 1).

Lueckisporites virkkiae, (Potonié and Klaus) Clarke, 1965 (Pl. VI, Fig. 3).

Mabuitasaccites ovatus, Bose and Kar, 1966 (Pl. IV, Fig. 11).

Marsupipollenites striatus, (Balme and Hennelly) Foster, 1979 (Pl. IV, Fig. 1).

Protohaploxylinus bharadwajii, Foster, 1979 (Pl. VI, Fig. 2).

Protohaploxylinus limpidus, (Balme and Hennelly) Balme and Playford, 1967 (Pl. VI, Fig. 6).

Striatoabieites multistriatus, (Balme and Hennelly) Hart, 1964 (Pl. V, Figs. 7 & 8).

Striatopodocarpites cancellatus, (Balme and Hennelly) Hart, 1963 (Pl. VI, Figs. 9 & 11).

Striatopodocarpites fusus, (Balme and Hennelly) Potonié, 1956 (Pl. VI, Fig. 5).

Striatopodocarpites raniganjensis, (Bharadwaj) Hart, 1964 (Pl. VI, Fig. 10).

Striomonosaccites brevis, Bose and Kar, 1966 (Pl. III, Fig. 3).

Striomonosaccites cicatricosus, Archangelsky and Gamarro, 1979 (Pl. III, Fig. 1).

Striomonosaccites triangularis, Maheshwari, 1967 (Pl. III, Fig. 2).

Tiwariaspis flavatus, Maheshwari and Kar, 1967 (Pl. III, Fig. 12).

Tiwariaspis gondwanensis, (Tiwari) Maheshwari and Kar, 1967 (Pl. IV, Fig. 4).

Trochosporites sp., (Pl. III, Fig. 5).

Tuberisaccites sp., (Pl. II, Fig. 4).

Vittatina costabilis, Wilson, 1962 (Pl. V, Fig. 3).

Vittatina subsaccata, Samoilovich, 1953 (Pl. V, Fig. 5).

Vittatina verrucosa, Tiwari, 1968 (Pl. V, Fig. 4).

Weylandites magnus, (Bose and Kar) Backhouse, 1991

Plate I

All x 1000 (bar = 20um)

- Figure 1. *Lophotriletes scotinus*, Segroves, 1970.
Figure 2. *Horriditriletes ramosus*, (Balme and Hennelly) Bharadwaj and Salujha, 196.
Figure 3. *Jayantisorites variabilis*, (Anderson) Backhouse, 1997.
Figure 4. *Pseudoreticulatispora pseudoreticulata*, (Balme and Hennelly) Bharadwaj and Srivastava, 1969.
Figure 5. *Punctatisporites gretensis*, Balme and Hennelly, 1956.
Figure 6. *Horriditriletes ramosus*, (Balme and Hennelly) Bharadwaj and Salujha, 1964.
Figure 7. *Densoisporites solidus*, Segroves, 1970.
Figure 8. *Verrucosisorites andersonii*, (Anderson) Backhouse, 1988.
Figure 9. *Leiotriletes virkkii*, Tiwari, 1965.
Figure 10. *Laevigatosporites colliensis*, (Bharadwaj and Hennelly) Venkatachala and Kar, 1968.
Figure 11. *Brazilea* sp.
Figure 12. *Kraeuselisporites niger*, Segroves, 1970.

Plate II

All x 1000 (bar = 20um)

- Figure 1. *Caheniasaccites ellipticus*, Maheshwari and Bose, 1968.
Figure 2. *Alisporites nuthallensis*, Clarke, 1965.
Figure 3. *Praecolpites ovatus*, (Anderson) Backhouse, 1991.
Figure 4. *Tuberisaccites* sp.
Figure 5. *Rugasaccites orbiculatus*, Lele and Maithy, 1969.
Figure 6. *Potonieisporites neglectus*, Potonié and Lele, 1959.
Figure 7. *Nuskoisporites triangularis*, Potonié and Lele, 1959.
Figure 8. *Falcisporites* sp.
Figure 9. *Plicatipollenites densus*, Srivastava, 1970.
Figure 10. *Caheniasaccites indicus*, Srivastava, 1970.
Figure 11. *Klausipollenites schaubergeri*, (Potonié and Klaus) Jansonius, 1962.
Figure 12. *Potonieisporites granulatus*, Bose and Kar, 1966.

Plate III

All x 1000 (bar = 20um)

- Figure 1. *Striomonosaccites cicatricosus*, Archangelsky and Gamero, 1979.
Figure 2. *Striomonosaccites triangularis*, Maheshwari, 1967.
Figure 3. *Striomonosaccites brevis*, Bose and Kar, 1966.
Figure 4. *Bascanisporites undosus*, Balme and Hennelly, 1956.
Figure 5. *Trochosporite* sp.
Figure 6. *Cycadopites cymbatus*, (Balme and Hennelly) Segroves, 1969.
Figure 7. *Crustaesporites globosus*, Leschik, 1956.
Figure 8. *Crustaesporites globosus*, Leschik, 1956.
Figure 9. *Weylandites magnus*, (Bose and Kar) Backhouse, 1991.
Figure 10. *Decussatisporites circularis*, Sinha, 1970.
Figure 11. *Distriamonocolpites ovalis*, Bharadwaj and Sinha, 1969.
Figure 12. *Tiwariasporis flavatus*, Maheshwari and Kar, 1967.

Plate IV

All x 1000 (bar = 20um)

- Figure 1. *Marsupipollenites striatus*, (Balme and Hennelly) Foster, 1979.
Figure 2. *Tiwariasporis flavatus*, Maheshwari and Kar, 1967.
Figure 3. *Doutakoffites elongatus*, Bose and Kar, 1966.
Figure 4. *Tiwariasporis gondwanensis*, (Tiwari) Maheshwari and Kar, 1967.
Figure 5. *Mabuitasaccites ovatus*, Bose and Kar, 1966.
Figure 6. *Costapollenites ellipticus*, Tschudy and Kosanke, 1966.
Figure 7. *Mabuitasaccites ovatus*, Bose and Kar, 1966.
Figure 8. *Fusacolpites ovatus*, Bose and Kar, 1966.
Figure 9. *Decussatisporites circularis*, Sinha, 1970.
Figure 10. *Boutakoffites elongatus*, Bose and Kar, 1966.
Figure 11. *Mabuitasaccites ovatus*, Bose and Kar, 1966.
Figure 12. *Fusacolpites fusus*, Bose and Kar, 1966.

(Pl. III, Fig. 9).

Acritarchs

Brazilea sp., (Pl. I, Fig. 11).

Gorgonisphaeridium sp., (Pl. V, Fig. 11).


Stratigraphic Palynology

This study was undertaken to determine the geological age of the formation, to make an interpretation of the


depositional site, and to reconstruct the palaeogeographic relationships of the Zagros Basin to Gondwana and Laurasia Supercontinents during the time interval represented by these strata. A total of 58 species are identified, including 46 pollen (32 genera), 10 spore (10 genera) and 2 acritarch (2 genera) species. The distribution of the taxa are shown on Table I and selected morphotype species are illustrated in Plates I-VI.

A detailed microscopic study of the palynological


Plate I




1



2




3




4




5




6




7




8




9



10




11




12

Plate II




9V


Plate III




1




2




3




4




5




6




7




8




9



10




11



12

Plate IV



99

samples of the Chal-i-Sheh Formation reveals that the base of this rock unit coincides with the appearance of abundant gymnospermous pollen taxa in all studied sections including:

Vittatina costabilis, *Costapollenites ellipticus*, *Hamiapollenites karrooensis*,

Potonieisporites granulatus, *Potonieisporites neglectus*, *Nuskoisporites triangularis*,

Plicatipollenites densus, *Plicatipollenites indicus*, *Rugasaccites orbiculatus*,

Klausipollenites schaubergeri, *Falcisporites flexuosa* and *Alisporites nuthalensis*

Diversity within this rock unit rapidly increases with the occurrence of other pollen species including:

Caheniasaccites ellipticus, *Caheniasaccites indicus*, *Fusacolpites fusus*, *Fusacolpites ovatus*,

Boutakoffites elongatus, *Mabuitasaccites ovatus*, *Cycadopites cymbatus*,

Marsupipollenites striatus, *Tiwariasporis flavatus*, *Tiwariasporis gondwanensis*,

Decussatisporites circularis, *Distriamonocolpites ovalis*, *Weylandites magnus*,

Bascanisporites undosus, *Vittatina subsaccata*, *Vittatina verrucosa*,

Walikalesaccites ellipticus, *Striatoabieites multistriatus*, *Striatopodocarpites fusus*,

Striatopodocarpites raniganjensis, *Striatopodocarpites cancellatus*, *Lueckisporites virkkiae*,

Protohaploxylinus limpidus, *Protohaploxylinus bharadwajii* and *Praecolpatites ovatus*.

In addition to the pollen taxa, there are several spore species in this rock unit including:

Laevigatosporites colliensis, *Leiotriletes virkkii*, *Punctatisporites gretensis*,

Lophotriletes scotinus, *Densoisporites solidus*, *Horriditriletes ramosus*,

Kraeuselisporites niger, *Pseudoreticulatispora pseudoreticulata*, *Verrucosisporites andersonii* and *Jayantisporites variabilis*

The relative percentage of each morphotype was calculated based on counts of two hundred grains per sample and the sum total resulted in 98% pollen, 1.5% acritarchs and scolecodonts.

Some of the components of this rock unit are found in the Early Permian Dorud Formation of northern Iran [19, 29] and the Early Permian of Turkey [1], but the majority of species of Chal-i-Sheh Formation are recorded in the Early Permian strata of Pakistan [68, 69, 70, 6], India [14, 68, 70, 59], Africa [32, 36, 15, 17, 2], Australia [55, 56, 31, 4] and a few species in the Early Permian sediments of the

United States of America [38, 66]. This indicates that the index species of Early Permian are:

Hamiapollenites karrooensis, *Vittatina costabilis*, *Vittatina subsaccata*, *Vittatina verrucosa*,

Costapollenites ellipticus, *Fusacolpites fusus*, *Fusacolpites ovatus*,

Boutakoffites elongatus, *Mabuitasaccites ovatus*, *Walikalesaccites ellipticus*,

Corisaccites alutas, *Nuskoisporites triangularis*, *Striomonosaccites brevis*,

Striomonosaccites triangularis, *Cycadopites cymbatus*, *Weylandites magnus*,

Potonieisporites granulatus, *Potonieisporites neglectus*, *Plicatipollenites densus*,

Plicatipollenites gondwanensis, *Decussatisporites circularis*, *Distriamonocolpites ovalis*,

Tiwariasporis flavatus, *Tiwariasporis gondwanensis*, *Striatopodocarpites fusus*,

Leiotriletes virkkii, *Verrucosisporites andersonii*, *Densoisporites solidus*, and other species are long-ranging during the Permian period (Table 1). Based on the above-mentioned species, an Early Permian (Sakmarian-Artinskian) age is suggested for the Chal-i-Sheh Formation in the Zagros Basin.

From the point of view of lithology and age relationship, the Chal-i-Sheh Formation is possibly equivalent to the Unayzah Formation in the southern part of the Persian Gulf (Saudi Arabia). The Upper Carboniferous age assignment to the Chal-i-Sheh Formation, based on the presence of the megafossil plant *Sigillaria persica* of Seward [58], is therefore rejected and the Carboniferous is represented by an hiatus in the Zagros Basin which possibly equates with the Hercynian Orogeny.

A small number of acritarch taxa and scolecodonts, which contribute 0.5 percent of the total assemblages, are associated with the miospore assemblages of the Chal-i-Sheh Formation. In all samples, the pollen diversity is high-masking the contributions of acritarchs, scolecodonts and spores. This phenomenon might be explained in terms of the close proximity of the depositional site to the parent plant source. Another interpretation would be that the Late Carboniferous Gondwana glaciation retreated and gave rise to relatively cold conditions at the beginning of the Permian period. This situation favored the gymnospermous plant community and inhibited the development of spore-bearing plants as well as acritarchs and marine annelid worms in the depositional environment. In summary, the Chal-i-Sheh Formation was deposited in a shallow marine environment.

Palaeogeography

The 58 morphotype species identified from the Early Permian sediments (Chal-i-Sheh Formation) of the Zagros

Plate V

All x 1000 (bar = 20um)

- Figure 1. *Hamiapollenites saccatus*, Wilson, 1962.
Figure 2. *Walikalesaccites ellipticus*, Bose and Kar, 1966.
Figure 3. *Vittatina costabilis*, Wilson, 1962.
Figure 4. *Vittatina verrucosa*, Tiwari, 1965.
Figure 5. *Vittatina subsaccata*, Samoilovitch, 1953.
Figure 6. *Vittatina subsaccata*, Samoilovitch, 1953.
Figure 7. *Striatoabieites multistriatus*, (Blame and Hennelly) Hart, 1964.
Figure 8. *Striatoabietites multistriatus*, (Balme and Hennelly) Hart, 1964.
Figure 9. *Vittatina costabilis*, Wilson, 1962.
Figure 10. *Corisaccites alutas*, Venkatachala and Kar, 1966.
Figure 11. *Gorgonisphaeridium* sp.
Figure 12. *Plicatipollenites indicus*, Lele, 1964.

Plate VI

All x 1000 (bar = 20 um)

- Figures 1&2. *Complexisporites polymorphus*, Jizba, 1962.
Figure 2. *Protohaploxypinus bharadwajii*, Foster, 1979.
Figure 3. *Lueckisporites virkkiae*, (Pottonié and Klaus) Clarke, 1965.
Figure 4. *Striatopodocarpites raniganjensis*, (Bharadwaj) Hart, 1964.
Figure 5. *Striatopodocarpites fusus*, (Balme and Hennelly) Pottonié, 1956.
Figure 6. *Protohaploxypinus limpidus*, (Balme and Hennelly) Balme and Playford, 1967.
Figure 7. *Hamiapollenites karrooensis*, (Hart) Hart, 1964.
Figure 8. *Hamiapollenites saccatus*, Wilson, 1962.
Figures 9 & 11. *Striatopodocarpites cancellatus*, (Balme and Hennelly) Hart, 1963.
Figure 10. *Striatopodocarpites raniganjensis*, (Bharadwaj) Hart, 1964.

Basin are considered to be sufficiently distinct and of potential value in reconstructing the palaeogeography of the Zagros Basin during the Early Permian. These species were compared with those of almost coeval strata in Africa [36, 15, 48, 17, 41, 67, 2], Australia [55, 56, 31, 4], India and Pakistan [14, 63, 69, 13, 6, 10], South America [22, 3, 45], Turkey [1], Europe and North America [38, 66, 21, 37]. This comparison indicates that from the 58 Early Permian species of the Zagros Basin, 40 species are common with Australia, 30 species with India and Pakistan, 47 species with Africa, 28 species with South America, 11 species with Europe and North America and three species with Turkey (Table 2). Comparison of the palynological data suggests a closer relation to those of Gondwanan countries than with those of Europe, North America and Turkey (Table 2). However, the Early Permian assemblage of the Zagros Basin appears to be most similar to those of African countries (Table 2). It would be reasonable, therefore, to suggest that the Zagros Basin and African Plate were not distant from one another and they were at about the same latitude along the southern shore of the Tethys sea during the Permian period.


Conclusions

The Early Permian sediments of the Zagros Basin are called the Chal-i-Sheh Formation, which is well-developed from SE-NW of this basin. The surface and subsurface samples of this rock unit were investigated for palynomorph entities. All samples contain well-preserved and abundant palynomorph taxa, which permit determination of the stratigraphic age of this formation.


A detailed microscopic study of the Chal-i-Sheh Formation reveals that the lower most part of the formation coincides with the appearance of abundant gymnospermous pollen taxa whose diversities increase toward the top of this rock unit. Among the miospore species of this rock unit, a few species have been recorded from the whole Permian period in other parts of the world, and most of them belong to the Early Permian (Table 1). Therefore, based on palynological data, there is a major hiatus between the Chal-i-Sheh Formation and older rock units and its magnitude increases from SE-NW of the Zagros Basin (Fig. 2). Likewise, the miospore species of this rock unit were compared with those of coeval strata elsewhere (Table 2). The comparison indicates that from 56 miospore species of Chal-i-Sheh Formation, 40 species are common with Australia, 47 species with Africa, 30 species with India and Pakistan, 28 species with South America, and three species with Turkey. Therefore, the Early Permian assemblage of Zagros Basin appears most similar to those of African countries. This similarity suggests that the Iranian platform and African plate were at about the same palaeolatitude along the southern shore of the Palaeo-Tethys Ocean during the Early Permian.

On the other hand, a cold limnic condition prevailed over the vegetation of the Zagros Basin during the Early Permian. This is indicated by the remnants of a Carboniferous forest (including *Sigillaria persica*) in the Chal-i-Sheh area and coal seams in Kuh-e-Gareh and Zard-Kuh area.


Plate V




1




2




3




4




5




6




7




8




9



10




11




12

lot


Plate VI




1




2




3




4




5




6




7




8




9



10



11



12

۱۰۳

Table I. Stratigraphic range chart of palynomorph taxa from the Early Permian strata in the Zagros Basin, of the Islamic Republic of Iran

LIST OF MORPHOTYPE SPECIES	U. Car bonif erous	Early Permian			Late Permian		
	Stephanian	Asselian	Sakmarian	Artinskian	Kungurian	Kazanian	Tatarian
<i>Hamiapollenites saccatus</i>							
<i>Horriditriletes ramosus</i>							
<i>Fusacolpites ovatus</i>							
<i>Fusacolpites fusus</i>							
<i>Pseudoreticulatispora pseudoreticulata</i>							
<i>Punctatisporites gretensis</i>							
<i>Potonieisporites neglectus</i>							
<i>Boutakoffites elongatus</i>							
<i>Mabuitasaccites ovatus</i>							
<i>Costapollenites ellipticus</i>							
<i>Walikalesaccites ellipticus</i>							
<i>Vittatina costabilis</i>							
<i>Vittatina verrucosa</i>							
<i>Nuskoisporites triangularis</i>							
<i>Plicatipollenites densus</i>							
<i>Potonieisporites granulatus</i>							
<i>Striomonosaccites cicatricosus</i>							
<i>Striomonosaccites triangularis</i>							
<i>Striomonosaccites brevis</i>							
<i>Cycadopites cymbatus</i>							
<i>Vittatina subsaccata</i>							
<i>Plicatipollenites indicus</i>							
<i>Complexisporites polymorphus</i>							
<i>Hamiapollenites karrooensis</i>							
<i>Protohaploxylinus bharadwajii</i>							
<i>Striatopodocarpites fusus</i>							
<i>Striatopodocarpites cancellatus</i>							
<i>Laevigatosporites colliensis</i>							
<i>Praecolpatites ovatus</i>							
<i>Crustasporites globosus</i>							
<i>Striatoabieites multistriatus</i>							
<i>Trochosporites sp.</i>							
<i>Verrucosisporites andersonii</i>							
<i>Leiotriletes virkkii</i>							
<i>Kraeuselisporites niger</i>							
<i>Marsupipollenites striatus</i>							
<i>Jayantisporites variabilis</i>							
<i>Corisaccites alutas</i>							
<i>Striatopodocarpites raniganjensis</i>							
<i>Protohaploxylinus limpidus</i>							
<i>Caheniasaccites ellipticus</i>							
<i>Tuberisaccites sp.</i>							
<i>Rugasaccites orbiculatus</i>							
<i>Caheniasaccites indicus</i>							
<i>Weylandites magnus</i>							
<i>Decussatisporites circularis</i>							
<i>Distriamocolpites ovalis</i>							
<i>Lophotriletes scotimus</i>							
<i>Densoisporites solidus</i>							
<i>Alisporites nuthallensis</i>							
<i>Klausipollenites schaubergeri</i>							
<i>Bascanisporites undosus</i>							
<i>Lueckisporites virkkiae</i>							
<i>Tiwariasporis flavatus</i>							
<i>Tiwariasporis gondwanensis</i>							

Table II. Palaeogeographic distribution of palynomorph taxa from the Early Permian Strata in the Zagros Basin of the Islamic Republic of Iran

LIST OF PALYNOMORPH TAXA FOUND IN THIS STUDY	Turkey	Europe & North America	South America	Africa	India & Pakistan	Australia
<i>Lophotriletes scotinus</i>	-	-	+	+	+	+
<i>Horridotriletes ramosus</i>	-	-	+	+	+	+
<i>Jayantispores variabilis</i>	+	-	+	+	-	+
<i>Pseudoreticulatispora pseudoreticulata</i>	-	-	+	+	+	+
<i>Punctatisporites gretensis</i>	+	-	+	+	+	+
<i>Densoisporites solidus</i>	-	-	-	+	-	+
<i>Ferrucosporites andersonii</i>	+	+	+	+	+	+
<i>Leiotriletes virkkii</i>	-	-	-	+	+	+
<i>Laevigatosporites colliensis</i>	-	-	+	+	+	+
<i>Kraeuselisporites niger</i>	-	-	-	+	+	+
<i>Caheniasaccites ellipticus</i>	-	-	+	+	+	+
<i>Alisporites nuthallensis</i>	-	-	-	+	+	+
<i>Praecolpites ovatus</i>	-	-	+	+	+	+
<i>Tuberisaccites sp.</i>	-	-	+	-	+	-
<i>Rugasaccites orbiculatus</i>	-	-	-	+	-	+
<i>Potonieisporites neglectus</i>	-	-	+	+	+	+
<i>Nuskoisporites triangularis</i>	-	-	-	-	+	-
<i>Falcisporites sp.</i>	-	-	-	-	-	-
<i>Plicatipollenites densus</i>	-	-	+	+	+	+
<i>Caheniasaccites indicus</i>	-	-	-	+	+	+
<i>Klausipollenites schaubergeri</i>	-	+	-	+	+	+
<i>Potonieisporites granulatus</i>	-	-	+	-	+	-
<i>Striomonosaccites cicatricosus</i>	-	-	+	-	-	-
<i>Striomonosaccites triangularis</i>	-	-	-	-	+	-
<i>Striomonosaccites brevis</i>	-	-	-	+	-	+
<i>Bascanisporites undosus</i>	-	-	+	-	-	+
<i>Trochosporites sp.</i>	-	+	-	+	-	-
<i>Cycadopites cymbatus</i>	-	-	+	+	+	+
<i>Crustaesporites globosus</i>	-	+	-	+	-	+
<i>Weylandites magnus</i>	-	-	+	+	+	+
<i>Decussatisporites circularis</i>	-	-	-	+	-	+
<i>Distriamonoalporites ovalis</i>	-	-	-	+	+	-
<i>Tiwariasporis flavatus</i>	-	-	+	+	+	+
<i>Marsupipollenites striatus</i>	-	-	+	+	-	+
<i>Boutakoffites elongatus</i>	-	-	-	+	+	+
<i>Tiwariasporis gondwanensis</i>	-	-	-	+	+	+
<i>Mabuitasaccites ovatus</i>	-	-	-	+	-	-
<i>Costapollenites ellipticus</i>	-	+	-	+	-	-
<i>Fusacolpites ovatus</i>	-	-	-	+	+	-
<i>Fusacolpites fusus</i>	-	-	-	+	+	-
<i>Hamiapollenites saccatus</i>	+	+	-	-	+	+
<i>Walikalesaccites ellipticus</i>	-	-	-	+	-	-
<i>Vittatina costabilis</i>	-	+	-	+	-	-
<i>Vittatina verrucosa</i>	-	-	-	+	+	-
<i>Vittatina subsaccata</i>	-	+	-	+	+	-
<i>Striatoabieites multistriatus</i>	-	-	+	+	-	+
<i>Corisaccites alutas</i>	-	-	+	+	+	+
<i>Plicatipollenites indicus</i>	-	-	+	+	+	+
<i>Complexisporites polymorphus</i>	-	+	-	-	+	-
<i>Protohaploxylinus bharadwajii</i>	-	-	+	+	+	+
<i>Lueckisporites virkkiae</i>	-	-	+	+	-	+
<i>Striatopodocarpites ramiganjensis</i>	-	-	-	+	+	+
<i>Striatopodocarpites fusus</i>	-	+	+	-	-	+
<i>Protohaploxylinus limpidus</i>	-	-	+	+	+	+
<i>Hamiapollenites karrooensis</i>	-	+	-	+	-	-
<i>Striatopodocarpites cancellatus</i>	-	+	+	+	+	+
<i>Brazilea sp.</i>	-	-	+	-	-	+
<i>Gorgonisphaeridium sp.</i>	-	-	+	-	+	+

100

Acknowledgements

I would like to express my sincere appreciation to Dr. Ralph E. Taggart, Professor of Botany and Plant Pathology, Department of Geological Sciences, Michigan State University, for his encouragement and assistance during all phases of this study. I would particularly like to express my appreciation to Dr. Aureal T. Cross, Professor Emeritus, Department of Geological Sciences, Michigan State University, for his constant advice and inspiration. I would like also to acknowledge, with gratitude, the Directory Board of the National Iranian Oil Company for the financial support given for all aspects of this research. Finally, I would like to express my thanks to my wife Fahimeh and my daughters Hedieh and Mona for their untiring encouragement and assistance.

References

- Akyol, E. Palynologie Permien Inferieur de Sariz (Kayseri) de Pamucak Yaylasi (Antalya` Turquie) et contamination Jurassique observee, due aux Ruisseaux "Pamucak" et "Goynuk". *Pollen et Spores*, 17, (1), 141-179, (1975).
- Anderson, J.M. The biostratigraphy of the Permian and Triassic. Part 3. A review of Gondwana palynology with particular reference to the northern Karroo Basin, South Africa. *Botanical Survey of South Africa, Memoir*, 41, 1-133, (1977).
- Archangelsky, S. and Gamarro, J. C. Palynologia del Paleozoico Superior en el Subsuelo de la Cuenca Chacoparaense, Republic Argentina. I. Estudio Sistemático de los paliniformes de tres perforaciones de la provincia de Cordoba. *Rev. Esp. Micropaleontol.*, 11, 417-478, (1979).
- Backhouse, J. Trilete spores from the Collie Basin, Western Australia. *Assoc. Aust. Palaeontol. Mem.*, 5, 53-72, (1988).
- Backhouse, J. Permian palynostratigraphy of the Collie Basin, Western Australia. *Rev. Palaeobot. Palynol.*, 67, (3/4), 237-314, (1991).
- Balme, B.E. Palynology of Permian and Triassic strata in the Salt Range and Sughar Range, West Pakistan, Stratigraphic Boundary Problems. Permian and Triassic of West Pakistan. *Univ. Kansas. Dept. Geol. Spec. Publ.*, 4, 306-453, (1970).
- Balme, B.E. and Hennelly, J.P.E. Trilete sporomorphs from Australian Permian sediments. *Aust. J. Bot.*, 3, (1), 240-260, (1956).
- Balme, B.E. and Playford, G. Late Permian plant microfossils from the Prince Charles Mountains Antarctica. *Rev. Micropaleontol.*, 10, (3), 192-197, (1967).
- Bharadwaj, D.C. The miospore genera in the coals of Raniganj Stage (Upper Permian) India. *Palaeobotanist*, 9, (1-2), 68-106, (1962).
- Bharadwaj, D.C. and Dwivedi, A. Spores-dispersae of the Barakar sediments from south Karanpura Coalfield, Bihar, India. *Ibid.*, 27, (1), 21-94, (1981).
- Bharadwaj, D.C. and Salujha, S.K. Sporological study of Seam VIII in Raniganj Coalfield, Bihar, India. Part I. Description of spores-dispersae. *Ibid.*, 12, (2), 181-215, (1964).
- Bharadwaj, D.C. and Sinha, V. Some new miospores from Early Gondwana coals. *J. Sen. Memorial*, 7-16, (1969).
- Bharadwaj, D.C. and Srivastava, S.C. Some new miospores from Barakar Stage, Early Gondwana, India. *Palaeobotanist*, 17, (2), 220-229, (1969).
- Bharadwaj, D.C. and Tiwari, R.S. On two monosaccate genera from Barakar Stage of India. *Ibid.*, 12, 181-215, (1964).
- Bose, M.N. and Kar, R.K. Palaeozoic spores dispersae from Congo. I. Kinda-Kalima and Walikale region. *Ann. Mus. R. Afr. Cent. Sci., Geol.*, in-8, 53: 1-238, (1966).
- Bose, M.N. and Kar, R.K. Palaeozoic spores dispersae from Zaire (Congo). XI. Assises glaciaires et periglaciaires from the Lukuga valley. *Mus. Roy. Afr. Cent. Terv. Belg. Ann. Ser. in-8, Sci. Geol.* 77, 3-13, (1976).
- Bose, M.N. and Maheshwari, H.K. Palaeozoic spores dispersae from Congo. VII: Coal measures near lake Tanganyika, south of Albertville. *Ibid.*, 60, 1-116, (1968).
- Broutin, J., Doubinger, J., Ousman El Hamet, M. and Lang, J. (Palynologie comparee du Permien nigerien (Afrique Occidentale) et peritethysien. Implications stratigraphiques et phytogeographique. *Rev. Palaeobot. Palynol.*, 66, (3/4), 243-261, (1990).
- Chateaufort, J. and Stampfli, G. Preliminary report on Permian palynology of Iran. *Proc. IV Int. Palynol. Conf. Lucknow, India (1976-1977)*, 2, 186-198, (1979).
- Clapham, Jr., W.B. Permian miospores from the Flowerpot Formation of western Oklahoma. *Micropaleontology*, 16, (1), 15-36, (1970).
- Clarke, R.F.A. British Permian saccate and monosaccate miospores. *Palaeontology*, 8, (2), 322-354, (1965).
- Cousminer, L. Permian spores from Apillapampa, Bolivia. *J. Palaeontol.*, 39, (6), 1097-1111, (1965).
- Foster, C.B. Permian plant microfossils from the Blair Athol Coal Measures, Central Queensland, Australia. *Palaeontographica*, Abt. B, 121-171, (1975).
- Foster, C.B. Permian plant microfossils of the Blair Athol Coal Measures, Baralaba Coal Measures and basal Rewan Formation of Queensland. *Queensl. Geol. Surv. Publ.*, 372. *Palaeontol. Pap.*, 45, 1-244, (1979).
- Foster, C.B. Spore-pollen assemblages of the Bowen Basin, Queensland (Australia): Their relationship to the Permian/Triassic boundary. *Rev. Palaeobot. Palynol.*, 36, (1-2), 165-183, (1982).
- Ghavidel-syooki, M. Palynostratigraphy and paleoecology of the Faraghan Formation, southeastern Iran. Ph.D. dissertation, Michigan State University, 279 pp. (unpublished), (1988).

27. Ghavidel-syooki, M. Palynological study of Palaeozoic sediments of the Chal-i-Sheh area southwestern Iran. *J. Sci. I. R. Iran.*, 4, (1), 32-46, (1993).
28. Ghavidel-syooki, M. Palynological study of Ordovician sediments and Faraghan Formation at Kuh-e-Surmeh, southern Iran. *Geosciences, Geological Survey of Iran, Scientific Quarterly Journal*, 3, (12), 28-35, (1994).
29. Ghavidel-syooki, M. Palynostratigraphy and palaeogeography of a Palaeozoic sequence in the Hassanakdar area, Central Alborz Range, northern Iran. *Rev. Palaeobot. palynol.*, 86, (1/2), 91-109, (1995).
30. Ghavidel-syooki, M. Biostratigraphy of acritarchs in the Paleozoic rock units in the Zagros Basin. CIMP Acritarch Subcommittee, Prague, Czech Republic, Abstr., p. 5, (1996).
31. Gilby, A.R. and Foster, C.B. Early Permian palynology of the Archaringa Basin south Australia. *Palaeontographica*, Abt. B, 109-191, (1988).
32. Hart, G. Microflora from the Ketewaka-Mchuchuma coalfield, Tanganyika, Tanganyika. *Geol. Surv. Bull.*, 36, 1-27, (1963).
33. Hart, G.F. A review of the classification and distribution of the Permian miospores: *Disaccate striatiti*. *C.R. 5th Congr. Int. Carb. Strat. Geol. Paris*, 3, 1171-1199, (1964).
- 33a. Harrison, J.V. The geology of some Salt Domes in Luristan. *Geol. Soc. London Quart. Jour.*, 86, 463-522, (1930).
34. James, G.A. and Wynd, J.G. Stratigraphic nomenclature of the Iranian Oil Consortium, Agreement area, (unpublished company report), (1965).
35. Jansonius, J. Palynology of Permian and Triassic Sediments, Peace River area, Western Canada. *Palaeontographica*, 110B, 35-98, (1962).
36. Jardine, S. Microflora des formations du Gabon Attribuees au Karroo. *Rev. Palaeobot. Palynol.*, 17, (1/2), 75-112, (1974).
37. Jerzykiewicz, J. Latest Carboniferous (Stephanian) and Early Permian (Autunian) palynological assemblages from the Intrasudetic Basin southwestern Poland. *Palynology*, 11, 117-131, (1987).
38. Jizba, K.M. Late Palaeozoic bisaccate pollen from the United States mid-continent area. *J. Paleontology*, 36, 871-887, (1962).
39. Kar, R.K. Palynology of the Barren Measures sequence from Jharia coalfield, Bihar, India. *Palaeobotanist*, 16, (2), 115-140, (1968).
40. Kar, R.K. Palynology of the north Karanpura Basin, Bihar, India-3. Raniganj exposure near Lungatoo Hazaribagh district. *Ibid.*, 16, (3), 275-282, (1968).
41. Kar, R.K. Palaeozoic spore dispersae from Congo. IX: Ombela and Lokandu regions (Lualaba river). *Mus. Roy. Afr. Cent. Terv. Belg. Ann. Serv. In-8, Sci. Geol.* 63, 84-105, (1969).
42. Kar, R.K. and Bose, M.N. Palaeozoic spore dispersae from Zaire (Congo). XII. Assise a couches de Houille from Greinerville region. *Ibid.*, 77, 23-113, (1976).
43. Lele, K.M. Studies in the Talchir Flora of India-2. Resolution of spore genus *Nuskosporites* Potonié and *Klaus*. *Palaeobotanist*, 12, (2), 168-147, (1964).
44. Leschik, G. Sporen aus dem Salton des Zechsteins von Neuhof (bei Fulda). *Palaeontographica*, 100, (B), 125-141, (1956).
45. Lindström, S. Early Late Permian palynostratigraphy and palaeo-biogeography of Vestfjella, Dronning Maudland, Antarctica. *Rev. Palaeobot. Palynol.*, 86, (1/2), 157-173, (1995).
46. Maheshwari, H.K. Studies in the Glossopteris Flora of India. 29: Miospore assemblage from the Early Gondwana exposures along Bansloi River in Rajmahal Hills Bihar. *Palaeobotanist*, 15, (3), 258-280, (1967).
47. Maheshwari, H.K. Palaeozoic spore dispersae from Congo. X. Microfossils from a cliff section at the confluence of Lufpa and Mushyashya rivers, south Katanga. *Mus. Roy. Afr. Cent. Terv. Belg. Ann. Ser. in-8, Sci. Geol.*, 63, 115-168, (1969).
48. Maheshwari, H.K. and Kar, M.N. *Tiwariasporis* Gen. nov. a new spore genus from the Permian of Congo and India. *Curr. Sci.*, 36, (14), 369-370, (1967).
49. Maheshwari, H.K. and Kar, M.N. Palaeozoic spore dispersae from Congo. VIII. The Kibamba river (Lukuga Coalfield area). *Mus. Roy. Afr. Cent. Terv. Belg. Ann. Ser. In-8. Sci. Geol.*, 63, 4-61, (1969).
50. Morgan, B.E. Two species of *Hamiapollenites* from the Wellington Formation of Oklahoma. *Micropaleontology*, 22, (1), 100-103, (1976).
51. Potonié, R. Synopsis der Gattungen der spore dispersae. II Teil. *Beih. Geol. Jb.*, 23, 1-103, (1956).
52. Potonié, R. and Klaus, W. Einige sporengattungen des alpien Salzgebirges. *Geol. Jb.*, 68, 517-546, (1954).
53. Potonié, R. and Lele, K.M. Studies in the Talchir Flora of India-1: Spore dispersae from the Talchir beds of south Rewa Gondwana Basin. *Palaeobotanist*, 8, (1-2), 22-36, (1961).
54. Samoilovich, S.R. Pollen und sporen der Permischen Ablagerungen von Tscherdin und Aktyubinsk im Vorural. *Arb. Erdol. Geol. Inst. U.S.S.R.*, 75, 5-57, (1953).
55. Segroves, K.L. Saccate plant microfossils from the Permian of Western Australia. *Grana Palynologica*, 9, (1-3), 174-227, (1969).
56. Segroves, K.L. Permian spores and pollen grains from the Perth Basin of Western Australia. *Ibid.*, 10, (1), 43-73, (1970).
57. Setudehnia, A.O. The Palaeozoic sequence at Zard-Kuh and Kuh-e-Dinar. *Iranian Petroleum Institute Bull.*, 6, 16-33, (1975).
58. Seward, A.C. A Persian *Sigillaria* Phil. *Trans. Roy. Soc.*

- London, Series 3, 221, B. 479, (1932).
59. Sinha, V. Spores-dispersae from Jhingurdah seam, Singrauli Coalfield (MP) India. *Palaeobotanist*, **19**, (2), 175-201, (1972).
 60. Srivastava, S.C. Mioflora investigation in some coal of Talchir Coalfield (Orissa) India. *Ibid.*, **18**, (2), 154-166, (1970).
 61. Stocklin, J. and Setudehnia, A.O. Stratigraphic Lexicon of Iran. *Geol. Survey Report No. 18*, 1-376, (1972).
 62. Szabo, F. and Kheradpir, A. Permian and Triassic stratigraphy Zagros Basin, south-west Iran. *J. Pet. Geol.*, **1**, (2), 57-82, (1978).
 63. Tiwari, R.S. Miospore assemblages in some coals of Barakar stage (Early Gondwana) of India. *Palaeobotanist*, **13**, 168-214, (1965).
 64. Tiwari, R.S. Palynological investigation of some coal seams in the Ib-river Coalfield (Orissa) India. *Ibid.*, **16**, (3), 222-242, (1968).
 65. Tiwari, R.S. and Awatar, R. Palynodating of Nidpur beds, Son Graben Madhya Pradesh. *Ibid.*, **38**, 105-121, (1989).
 66. Tschudy, R.H. and Kosanke, R.M. Early Permian vesiculate pollen from Texas. *Ibid.*, **15**, (1-2), 59-71, (1966).
 67. Utting, J. Pollen and spore assemblages in the Luwumbu Coal Formation (Early Karroo) of the north Luangwa Valley Zambia and their biostratigraphic significance. *Rev. Palaeobot. Palynol.*, **21**, 295-315, (1976).
 68. Venkatachala, B.S. and Kar, R.K. Schizopollis Venkatachala and Kar, a new pollen genus from the Permian of north Karanpura Coalfield, Bihar, India. *Grana Palynologia*, **5**, (3), 415-424, (1964).
 69. Venkatachala, B.S. and Kar, R.K. *Corisaccites* Gen. nov. A new saccate pollen genus from the Permian of Salt Range, West Pakistan. *Palaeobotanist*, **15**, (1-2), 107-109, (1966).
 70. Venkatachala, B.S. and Kar, R.K. Palynology of the north Karanpura Basin, Bihar, India-2. Barakar exposure near Lungatoo, Hazaribagh district. *Ibid.*, **16**, (3), 258-268, (1968c).
 71. Venkatachala, B.S. and Kar, R.K. Palynology of the Kathwai Shales, Salt Range, west Pakistan. Shales 25ft. above the Talchir Boulder Beds. *Ibid.*, **16**, (2), 156-166, (1968b).
 72. Venkatachala, B.S. and Kar, R.K. Palynology of the Karanpura sedimentary basin, Bihar, India-1. Barakar Stage at Badam. *Ibid.*, **16**, (1), 56-90, (1968a).
 73. Vijaya, R. Evolutionary pattern of striations and taeniae in the Indian Gondwana saccate pollen. *Ibid.*, **38**, 83-91, (1990).
 74. Warrington, G. and Scrivener, R.C. The Permian of Devon, England. *Rev. Palaeobot. Palynol.*, **66**, (3/4), 263-272, (1990).
 75. Wilson, L.R. Permian plant microfossils from the Flowerpot Formation, Greer Country, Oklahoma. *Okla. Geol. Surv. Circ.*, **49**, 5-502, (1962).